

Postpartum depression in adolescent mothers before and during COVID-19 and the role of self-esteem, maternal self-efficacy, and social support

Pamela Patiño,¹ María Asunción Lara,² Corina Benjet,³ Asunción Alvarez-del Río,⁴ Feliciano Bartolo Solís⁵

¹ Programa de Maestría y Doctorado en Ciencias Médicas, Odontológicas y de la Salud, Facultad de Medicina, Universidad Nacional Autónoma de México, Ciudad de México, México.

² Dirección de Investigaciones Epidemiológicas y Psicosociales, Departamento de Estudios Psicosociales en Poblaciones Especiales, Instituto Nacional de Psiquiatría Ramón de la Fuente Muñiz, Ciudad de México, México.

³ Dirección de Investigaciones Epidemiológicas y Psicosociales, Centro de Investigación en Salud Mental Global, Instituto Nacional de Psiquiatría Ramón de la Fuente Muñiz, Ciudad de México, México.

⁴ Departamento de Psiquiatría y Salud Mental, Facultad de Medicina, Universidad Nacional Autónoma de México, Ciudad de México, México.

⁵ Dirección de Atención Médica, Servicios de Salud Pública de la Ciudad de México, México.

Correspondence:

María Asunción Lara Cantú
Dirección de Investigaciones Epidemiológicas y Psicosociales. Instituto Nacional de Psiquiatría Ramón de la Fuente Muñiz.

Calz. México-Xochimilco 101,
Col. San Lorenzo Huipulco,
Alcaldía. Tlalpan, C.P. 14370,
Ciudad de México, México.
Phone: +52 (55) 4160-5170
Fax: +52 (55) 5513-3446
Email: laracan@imp.edu.mx

Received: 13 July 2023

Accepted: 15 August 2023

Citation:

Patiño, P., Lara, M. A., Benjet, C., Alvarez del Río, A., & Bartolo Solís, F. (2024). Postpartum depression in adolescent mothers before and during COVID-19 and the role of self-esteem, maternal self-efficacy, and social support. *Salud Mental*, 47(1), 23-33.

DOI: 10.17711/SM.0185-3325.2024.004



ABSTRACT

Introduction. Although the COVID-19 pandemic negatively impacted the mental health of vulnerable populations, such as adolescent mothers, very few studies have documented the prevalence of postpartum depression (PPD) in this population. **Objective.** a) Determine the frequency of PPD (Edinburgh Postnatal Depression Scale [EPDS] ≥ 9) in adolescent mothers before (AM-BP) and during (AM-DP) the pandemic, b) Examine psychosocial factors (self-esteem, maternal efficacy, social support, depression and anxiety in pregnancy, planned and wanted pregnancy) in AM-BP and AM-DP, and c) Determine whether being an AM-DP was a significant factor for experiencing PPD (EPDS ≥ 9). **Method.** Cross sectional study. Subjects: Forty-one AM-BP recruited at Health Centers and interviewed face to face and forty-one AM-DP surveyed online. **Results.** PPD (EPDS ≥ 9) was 42% ($p = .001$) more frequent in AM-DP. The groups differed significantly in all psychosocial factors, with AM-DP faring worse. Unadjusted regressions showed that being an AM-DP, having lower maternal efficacy and self-esteem, greater dissatisfaction with social support, and depression and/or anxiety in pregnancy increased PPD (EPDS ≥ 9). Adjusted multiple analysis indicated that lower self-esteem was the only factor to maintain its association with PPD (EPDS ≥ 9 ; $p = .017$). **Discussion and conclusion.** The pandemic negatively affected PPD (EPDS ≥ 9) and psychosocial factors in AM-DP, as compared to AM-BP, with self-esteem being the main factor associated with PPD (EPDS ≥ 9). In situations of extreme stress as happened in the pandemic, the mental health of adolescent mothers should be prioritized to prevent negative effects such as PPD. PPD preventive and treatment interventions should consider strengthening self-esteem.

Keywords: Adolescents, postpartum depression, maternal self-efficacy, self-esteem, social support, COVID-19.

RESUMEN

Introducción. La pandemia por COVID-19 tuvo un impacto negativo en la salud mental de poblaciones vulnerables, como las madres adolescentes, no obstante, escasos estudios documentaron la prevalencia de depresión posparto (DPP) en esta población. **Objetivo.** a) Conocer la frecuencia de DPP (Escala Edinburgh para la Depresión Postnatal [EPDS] ≥ 9) en madres adolescentes antes de la pandemia (MA-AP) y durante la pandemia (MA-DP), b) Examinar algunos factores psicosociales (autoestima, eficacia materna, apoyo social, depresión y ansiedad en el embarazo, embarazo planeado y deseado) en MA-AP y MA-DP, y, c) Analizar si ser MA-DP, fue un factor significativo para experimentar DPP (EPDS ≥ 9). **Método.** Estudio transversal. Participantes: 41 MA-AP captadas en Centros de Salud y 41 MA-DP encuestadas en línea. **Resultados.** La DPP (EPDS ≥ 9) fue 42% ($p = .001$) más frecuente en las MA-DP. Los grupos difirieron significativamente en todos los factores psicosociales, en detrimento de las MA-DP. Las regresiones no ajustadas mostraron que ser MA-DP, tener menor eficacia materna y autoestima, mayor insatisfacción con el apoyo social, y depresión y/o ansiedad en el embarazo incrementaron la DPP (EPDS ≥ 9). El análisis múltiple ajustado indicó que una menor autoestima fue el único factor que mantuvo su asociación con DPP (EPDS ≥ 9 ; $p = .017$). **Discusión y conclusión.** La pandemia tuvo un efecto negativo en la DPP (EPDS ≥ 9) y en factores psicosociales en MA-DP; la autoestima fue el principal factor asociado a la misma. Ante situaciones de estrés extremo, la salud mental de madres adolescentes debería ser prioritaria para prevenir efectos negativos como la DPP. Intervenciones preventivas y de tratamiento de DPP deben fortalecer la autoestima.

Palabras clave: Adolescentes, depresión posparto, eficacia materna, autoestima, apoyo social, COVID-19.

INTRODUCTION

The pandemic intensified a range of social and health problems (WHO, 2022; Edward & Mobarak, 2022), particularly affecting women (Burki, 2020). According to experts, unmet needs for contraceptives and safe abortion, lack of reproductive health programs, domestic sexual violence, poverty, and school dropout rates rose during the pandemic (NU - ECLAC, 2020; Borg Xuereb et al., 2023), contributing to the increase in the adolescent fertility rate (NU - ECLAC, 2020), which rose by 30% in Mexico (UNFPA, 2023).

Adolescents who experienced pregnancy and motherhood during the pandemic were forced to cope with adversity and intense emotions; fear of having loved ones with COVID-19, stress due to unemployment and financial difficulties (Astle et al., 2021; Smiley et al., 2021; Merriman et al., 2023), and uncertainty due to changes in their prenatal care and birth plans (Astle et al., 2021; Moltrecht et al., 2022). Many reported feeling unprepared to stimulate their children's development because they had been unable to take advantage of prenatal classes and contact with adults who would otherwise have provided guidance (Moltrecht et al., 2022).

Some adolescents reported feeling depressed due to the strict lockdown they observed out of fear of the unknown consequences of COVID-19 on pre-existing medical conditions, their pregnancy, or their newborns. Conversely, those who interrupted social distancing to receive support from their families reported experiencing anxiety and guilt (Moltrecht et al., 2022; Merriman et al., 2023). Adolescent mothers who parented their babies without support from their extended families and or partners—who were usually the providers—experienced constant frustration (Merriman et al., 2023). Those who continued their studies during the pandemic found it overwhelming to balance them with motherhood. Many were forced to interrupt or drop out of their studies because they lacked support to care for their children (Smiley et al., 2021).

In 2020, the adversities and intense emotions caused by the pandemic were expected to contribute to mental health problems (Shigemura et al., 2020), and a recent meta-review showed that depression and anxiety increased in adolescents, pregnant and postpartum women and people hospitalized with COVID-19 (Bower et al., 2023). Three research projects were found on the mental health of adolescent mothers during the pandemic. The first reported lower parental stress and postpartum depression (PPD) during the pandemic than before it started, attributed to a life skills program (LSP) and continuous school contact (Astle et al., 2021). The second research project reported a low level of PPD associated with good family support in two groups of adolescent mothers: under lockdown vs. not under lockdown (Matei et al., 2021), while the third found a higher proportion of PPD and lower social support among

adolescent than adult mothers in the group (Sangsawang & Sangsawang, 2023).

These research projects during the pandemic coincide with studies prior to this one showing that adolescent and adult mothers who receive adequate postpartum support are less likely to experience PPD and that the protective effect is greater for adolescents (deCastro et al., 2011; Kim et al., 2014).

Self-esteem and confidence in one's ability to solve problems—associated with parenting or otherwise—increases the use of effective coping strategies, maintaining hope and commitment under challenging circumstances (Bandura, 1994; Lazarus & Folkman, 1987). These characteristics can play an important role in coping with the stressors of motherhood, the pandemic, or both. Research on adolescent mothers reports that low self-esteem (Birkeland et al., 2005; Ramos-Marcuse et al., 2009) and a negative perception of their own abilities to foster the development of infants and overcome parenting difficulties—low perception of maternal efficacy—are associated with PPD (Lara et al., 2017; Léniz-Maturana et al., 2022). Sangsawang & Sangsawang (2023) observed that the maternal efficacy of adolescent girls decreased due to social distancing, which was associated with PPD. Before the pandemic, children of adolescent mothers with lower maternal efficacy were found to have lower emotional self-regulation (Léniz-Maturana et al., 2022).

In addition to the difficulties associated with COVID-19, even under normal circumstances, adolescent mothers are more likely to have obstetric complications, and premature babies (< 37 weeks of gestation or gw) with low birth weight (< 2500g; Ganchimeg et al., 2014). In some contexts, adolescent mothers are stigmatized, especially if they do not have a partner, which reduces social support (Merriman et al., 2023; Moltrecht et al., 2022), except from their mothers, who tend to support them (Alvarez-Nieto et al., 2014). Moreover, many fail to overcome the cycle of poverty due to educational lag (UNFPA, 2020). The disadvantages of adolescent mothers contribute to depression, which can affect their parenting and increase the risk of behavioral problems in their children (Hodgkinson et al., 2014). In Mexico, between 16.05% (deCastro et al., 2011) and 33.3% (Patiño, 2016) of adolescents have been found to have PPD.

In short, previous studies focused on the challenges and emotions faced by adolescent mothers during the pandemic, tending to overlook the detection and analysis of factors associated with PPD and other mental health problems in this population. Only one gathered data on PPD before the pandemic and during the first months of lockdown (Astle et al., 2021). No studies were conducted in Mexico. The data available in this country show that, during the COVID-19 health crisis, pregnant adults suffered stress (32%) and depression (17.5%; Medina-Jimenez et al., 2022), greater stress was associated with higher levels of depression and

social support was associated with less depression (Rivera-Rivera et al., 2021). Likewise, it was observed that experiencing pregnancy and postpartum during the pandemic was overwhelming due to the intense fear of contagion and the multiple demanding situations (Lara et al., 2023). Having data on the effects of the COVID-19 pandemic on the mental health and well-being of postpartum adolescent mothers will fill a gap in research, both in Mexico and in other countries, and serve as the basis for PPD prevention and management in this population in future crises.

Objectives

a) Determine the frequency of PPD (EPDS ≥ 9) in adolescent mothers before (AM-BP) and during the pandemic (AM-DP), b) Examine certain psychosocial factors (self-esteem, maternal efficacy, social support, depression and pregnancy anxiety, planned and desired pregnancy) in AM-BP and AM-DP, and c) Analyze whether belonging to the AM-DP group was a significant factor for experiencing PPD (EPDS ≥ 9) when adjusting for other sociodemographic, obstetric, and psychosocial factors.

METHOD

Design of the study/Places

A cross-sectional, comparative, correlational study was conducted with two groups of adolescent mothers. The first was surveyed from July 2019 to March 2020 at two Primary Health Care Centers (HC) in Mexico City (CDMX): adolescent mothers before the pandemic (AM-BP). Recruitment at HC was halted due to lockdown. The second group was surveyed online from March 8 to July 8, 2021: adolescent mothers during the pandemic (AM-DP).

Participants

The inclusion criteria were being a first-time mother, aged 14 to 19, with a baby aged two to nine months and living in Mexico City. In addition, AM-DPs were required to have an email address so they could access the online questionnaire. The exclusion criterion was for the mother to have reported having one of the following situations: 1) health problems during childbirth or postpartum requiring hospitalization for a week or longer or that caused pain and made it difficult for her to perform her everyday activities, 2) health problems in the past month unrelated to childbirth or postpartum causing hospitalization, 3) having a baby with health problems requiring hospitalization, or 4) having had very low birth weight (≤ 1.5 kg) or being born before 32 weeks' gestation.

The sample of 82 teenage mothers was non-probabilistic and included 41 AM-BP and 41 AM-DP. The groups

were matched by mother and baby ages to ensure representativeness of ages. They were also matched by whether the adolescents were in a relationship with the baby's father, because the majority of the AM-BP had a relationship with the fathers of their babies ($n = 31, 75.6\%$).

Measurements

Sociodemographic data: age of the mother (years) and baby (months), educational attainment, whether the mother attended school during pregnancy and postpartum, whether she worked postpartum, marital status, whether she had a relationship with the baby's father postpartum. Socioeconomic status (SES) was evaluated using the Mexican Association of Market Intelligence and Opinion Agencies Index (AMAI, 2020), comprising seven levels: A/B high (≥ 205 points), C+ or upper-middle (166-204 points), C or middle (136-165 points), C- or lower-middle (112-135 points), D+ low (90-111 points), D poverty (48-89 points), and E extreme poverty (0-47 points; AMAI, 2020; López-Romo, 2020).

Obstetric data: Birthweight (in grams) and number of weeks of gestation at the time of delivery.

The Edinburgh Postnatal Depression Scale (EPDS; Cox et al., 1987) was used to identify mothers with a high probability of PPD rather than to establish a clinical diagnosis. It consists of ten statements with a Likert-type response (0 to 3), with the total score being calculated by adding the scores of each item. The higher the score, the greater the likelihood of PPD. Although research on the optimal cut-off point of the EPDS to identify adolescents with a high probability of depression during pregnancy and postpartum is incipient, it has been recommended to use the score ≥ 9 (Barassi & Grealish, 2022). In the validation of the instrument with low-income Mexican pregnant adolescents the optimal cut-off point (8/9), adequate sensitivity (70.4%), and specificity (84.9%; Alvarado-Esquivel et al., 2014) were obtained. With primiparous adolescent mothers in postpartum, the cut-off point ≥ 9 had a sensitivity and specificity of 90% (Venkatesh et al., 2014), which is why this cut-off point was used in this study.

The Postpartum Depression Predictors Inventory (PDI-R; Beck, 2001; Records et al., 2007), was validated in Mexico (Ibarra-Yruegas et al., 2016) and administered to adolescent mothers (Lara et al., 2017; Patiño, 2016). Several features from the inventory were used to evaluate the psychosocial factors described below. Social support: This scale comprises twelve questions evaluating whether a woman has adequate support (from family, partner, and friends). Questions are answered with yes (0) or no (1). The total score is the sum of each item: the higher the score, the greater the participant's dissatisfaction with the support she received. Reliability of the social support scale in this study was higher than $\alpha = .70$ in AM-BP and AM-DP. 2) Depres-

sion and anxiety in pregnancy. One item was used to evaluate the perception of depression and another to evaluate the perception of anxiety during pregnancy, both of which are a retrospective measurement. The response options for each item are yes (1) or no (0). 3) Planned and desired pregnancy. One item was used to determine whether the pregnancy was planned and another to determine whether it was wanted. The response options for both questions are yes (0) or no (1).

Maternal self-esteem and efficacy were evaluated using the [Rosenberg Self-Esteem Scale \(1965\)](#) and the Parental Evaluation Scale (PES; [Farkas-Klein, 2008](#)). The Rosenberg scale comprises 10 statements with Likert-type responses (1 to 4). The total score is the sum of the items; the more the total score increases, the higher the level of self-esteem. Reliability is $\alpha = .86$ ([Ramos-Marcuse et al., 2009](#)). The PES, designed to evaluate overall self-perception of maternal efficacy, has 10 statements answered from 0 to 10. As the total score increases, the perception of maternal efficacy rises. Reliability is $\alpha = .77$ ([Léniz-Maturana et al., 2022](#)).

Procedure

- AM-BP. Those who attended the immunization appointment with their babies at the HC were invited to answer the survey. The survey was administered if they met the selection criteria and signed the informed consent form (IC). Any doubts the participants had were clarified before they signed the IC. The author and two trained psychology interns administered the surveys.
- AM-DP. These adolescent mothers were recruited through an online survey developed with the Google forms tool. The survey was published on the Facebook page “Help for Depression” (Administered by a research project of the Ramón de la Fuente Muñiz National Institute of Psychiatry [INPRFM]). Paid advertising was used to disseminate the survey on Facebook among the population of interest for the study, and the scope criteria were defined by sex, age, topics of interest, and location (women ages 14 to 19, with an interest in motherhood issues, postpartum and health and residing in Mexico City). In the first part of the survey, the justification and objectives of the study were presented. Those interested registered their email to see the IC and those who accepted the research conditions accessed the filter questions (selection criteria). One hundred and fourteen responses were received and 41 were finally included (Figure 1).

Statistical analyses

Descriptive statistics were obtained, and comparison analyses conducted of the groups (AM-BP and AM-DP) for

sociodemographic, obstetric, psychosocial, and PPD variables ($EPDS \geq 9$). X^2 (categorical variables), Student's t for independent samples (quantitative variables meeting the assumptions of normality and homogeneity of variances), and Mann-Whitney U (quantitative variables that did not meet the assumption of normality) tests were performed. The assumptions of normality and homogeneity were evaluated using the Kolmogorov-Smirnov and Levene tests. Differences between the groups of paired variables were not analyzed.

The following analyses were conducted to determine whether there was a greater probability of PPD ($EPDS \geq 9$) (dependent variable) among AM-DP than in AM-BP when adjusting for other sociodemographic, obstetric, and psychosocial factors (independent variables): simple logistic regression models to evaluate the association between the groups (AM-BP vs. AM-DP) and sociodemographic, obstetric and psychosocial factors (independent variables), and PPD ($EPDS \geq 9$; dependent variable). A multiple logistic regression model (adjusted) was built using significant variables from previous regressions.

A significance of $p < .05$ was used for the difference and association tests. Analyses were performed using SPSS Version 25 ([International Business Machines \[IBM\], 2017](#)).

Ethical considerations

This research was approved by the INPRFM Research Ethics Committee (Initial CEI/C/096/2018. Addendum CEI/C/005/2021). All participants gave their consent prior to their inclusion in the study. In return for their participation, they were given a pamphlet on PPD and a directory of psychological, legal, and sexual health advice.




Figure 1. Diagram of adolescent mothers recruited during the pandemic (AM-DP) who answered the online survey.

RESULTS

Sociodemographic and obstetric factors

As can be seen from Table 1, the average age of mothers was 17.7 years ($SD = 1.3$) and the average age of babies 4.7 months ($SD = 2.2$), with 45.1% having middle/lower middle SES and 50% having completed their basic education. There were no differences between groups in regard to sociodemographic or obstetric factors.

PPD (EPDS ≥ 9) and psychosocial factors

As shown in Table 2, 26.8% of AM-BP reported PPD (EPDS ≥ 9) compared to 68.35% of AM-DP, in other words, PPD (EPDS ≥ 9) was 42% higher in MA-DP ($p = .001$). In regard to psychosocial factors, it was found that AM-DP had significantly lower scores in maternal efficacy ($p = .001$) and self-esteem ($p = .008$), as well as greater dissatisfaction with their overall social support ($p = .030$) and support from their families ($p = .022$) than AM-BP. Furthermore, AM-DPs

reported feeling depressed (82.9%; $p = .001$) and/or anxious (70.7%; $p = .014$) during pregnancy more frequently than AM-BPs. No differences were found between the groups in terms of wanted and planned pregnancies.

Association between groups (AM-BP vs AM-DP), sociodemographic, obstetric, and psychosocial factors and PPD (EPDS ≥ 9)

Table 3 shows the logistic regression models (simple) used to analyze the association between the groups (AM-BP vs. AM-DP) and sociodemographic, obstetric, and psychosocial factors and PPD (EPDS ≥ 9). The results show that belonging to the AM-DP group ($p = .001$), having lower maternal efficacy ($p = .001$), lower self-esteem ($p = .001$), and greater dissatisfaction with overall social support ($p = .001$), their partners ($p = .006$) and family ($p = .013$), and presenting with depression ($p = .001$) and anxiety ($p = .013$) during pregnancy increased the probability of PPD (EPDS ≥ 9). Table 4 shows the multiple logistic regression model, adjusted for factors such as the group to which the

Table 1
Descriptive analyses and comparison of sociodemographic and obstetric factors

	AM-BP (n = 41)	AM-DP (n = 41)	Total (N = 82)	p
Sociodemographic data				
Mother's age (years)	17.7 ($SD = 1.3$)	17.7 ($SD = 1.3$)	17.7 ($SD = 1.3$)	NA
Baby's age (months)	4.7 ($SD = 2.2$)	4.7 ($SD = 2.2$)	4.7 ($SD = 2.2$)	NA
SES				
A/B and C+ (High /upper middle)	10 (24.4%)	8 (19.5%)	18 (22.0%)	
C and C - (Middle/lower-middle)	21 (51.2%)	16 (39.0%)	37 (45.1%)	.258 ^a
D+ D and E (Low level/poverty)	10 (24.4%)	17 (41.5%)	27 (32.9%)	
Educational Attainment				
Basic	21 (51.2%)	20 (48.8%)	41 (50%)	.825 ^a
Middle school	20 (48.8%)	21 (51.2%)	41 (50%)	
Attended school at the time they became pregnant				
Yes	16 (39.0%)	19 (46.3%)	35 (42.7%)	.503 ^a
No	25 (61.0%)	22 (53.7%)	47 (57.3%)	
Attended school postpartum				
Yes	9 (22.9%)	13 (31.7%)	22 (26.8%)	.319 ^a
No	32 (78.0%)	28 (68.3%)	60 (73.2%)	
Marital status				
Married/Living together	28 (68.3%)	29 (70.7%)	57 (69.5%)	.810 ^a
Single/Separated	13 (31.7%)	12 (29.3%)	25 (30.5%)	
In a relationship with the baby's father				
Yes	31 (75.6%)	31 (75.6%)	62 (75.6%)	NA
No	10 (24.4%)	10 (24.4%)	20 (24.4%)	
Work (at the time of the survey)				
Yes	6 (14.6%)	5 (12.2%)	11 (13.4%)	.746 ^a
No	35 (85.4%)	36 (87.8%)	71 (86.6%)	
Obstetrics				
Gestation weeks at delivery	39.1 ($SD = 1.4$)	38.6 ($SD = 1.7$)	38.8 ($SD = 1.6$)	.343 ^b
Birthweight (g)	2984.7 ($SD = 456.2$)	2961.2 ($SD = 402.1$)	2972.9 ($SD = 427.5$)	.805 ^c

Notes: AM-BP = Adolescent mothers before the pandemic; AM-DP = Adolescent mothers during the pandemic; SES = Socioeconomic Status; GW = Gestation weeks at delivery.
Test performed: ^a X²; ^b U Mann Whitney; ^c t test.

Table 2
Descriptive analyses and group comparison: PPD (EPDS \geq 9) and psychosocial factors

	AM-BP (n = 41)	AM-DP (n = 41)	Total (N = 82)	p
PPD				
Yes (EPDS \geq 9)	11 (26.8%)	28 (68.35%)	39 (47.6%)	.001 ^a
No (EPDS < 9)	30 (73.2%)	13 (31.7%)	43 (52.4%)	
Maternal efficacy R [0-10]	7.6 (SD 1.6)	6.2 (SD 1.8)	6.9 (SD 1.8)	.001 ^c
Self-esteem R [10-40]	32.0 (SD 4.8)	28.7 (SD 6.1)	30.4 (SD 5.7)	.008 ^b
Dissatisfaction with overall social support and by type of source				
Overall or total [†] R [0-12]	3.8 (SD = 2.4)	5.1 (SD = 2.7)	4.4 (SD = 2.6)	.030 ^b
Partner R [0-4]	1.2 (SD = 1.7)	1.6 (SD = 1.5)	1.4 (SD = 1.6)	.189 ^b
Family R [0-4]	.34 (SD = .79)	1.0 (SD = 1.3)	.6 (SD = 1.1)	.022 ^b
Friends R [0-4]	2.2 (SD = 1.6)	2.5 (SD = 1.6)	2.4 (SD = 1.6)	.321 ^b
Depression during pregnancy				
Yes	13 (31.7%)	34 (82.9%)	47 (57.4%)	.001 ^a
No	28 (68.3%)	7 (17.1%)	34 (42.7%)	
Anxiety during pregnancy				
Yes	18 (43.9%)	29 (70.7%)	47 (57.4%)	.014 ^a
No	23 (56.1%)	12 (29.3%)	35 (42.7%)	
Planned pregnancy				
Yes	9 (22.2%)	14 (34.1%)	23 (28.0%)	.219 ^a
No	32 (78.0%)	27 (65.9%)	59 (72.0%)	
Wanted pregnancy				
Yes	28 (68.3%)	28 (68.3%)	56 (68.3%)	1.00 ^a
No	13 (31.7%)	13 (31.7%)	26 (31.7%)	

Notes: AM-BP = Adolescent mothers before the pandemic; AM-DP = Adolescent mothers during the pandemic; R = Response range. Test performed: ^a χ^2 ; ^b U Mann Whitney; ^c t test.

[†] The higher the score, the greater the dissatisfaction with social support.

mother belongs (AM-BP and AM-DP), maternal efficacy, self-esteem, dissatisfaction with overall social support, depression, and anxiety during pregnancy. The results indicate that the AM-DP group was no longer significant for PPD (EPDS \geq 9) when other factors were adjusted for. In this model, self-esteem was the only prevailing factor associated with PPD (EPDS \geq 9; $p = .017$). The data fit well with the multiple logistic regression model ($\chi^2 = 55.995$; $p = .001$).

DISCUSSION AND CONCLUSION

The first objective of the study was to determine the frequency of PPD (EPDS \geq 9) in AM-BP and AM-DP. The results show that PPD (EPDS \geq 9) was 41.5% more frequent in AM-DP than in AM-BP, which was significant given that the groups had similar sociodemographic and obstetric features. This increase may reflect the stressful situations young mothers faced: fear of having loved ones with COVID-19, financial problems, and fear of interacting with other people (Astle et al., 2021; Smiley et al., 2021; Merriman et al., 2023).

A comparison of the PPD (EPDS \geq 9) of the AM-DP (68.35%, EPDS \geq 9) with similar research showed that it was higher than that of other adolescent mothers who also experienced their postpartum during the pandemic (36%,

EPDS \geq 13; Sangsawang & Sangsawang, 2023). These differences can be attributed to the use of different cut-off points, the fact that this study was based on an online survey, and the context and risk factors to which Mexican adolescent mothers were exposed. In regard to context, data in Sangsawang & Sangsawang (2023) were gathered during lockdown, whereas those for this article were collected between March 8 and July 8, 2021, following a spike in infections and deaths due to the pandemic in Mexico. During this period, it was confirmed that although the country had exceeded the worst-case scenario threefold, compulsory lockdown was not restored (Sánchez-Tanquer et al., 2021).

The analysis of sociodemographic, obstetric, and psychosocial factors before and during the pandemic showed differences in all psychosocial factors, with AM-DP displaying lower self-esteem, lower maternal efficacy, greater dissatisfaction with overall social and family support and a history of anxiety and depression during pregnancy. These results coincide with previous research reporting that, the greater the perception of stress due to the pandemic, the lower the parenting effectiveness-in adults (Xue et al., 2021; Gniewosz, 2022) and the lower the self-esteem of adolescents (Goto et al., 2022). They are also consistent with research showing a lack of support from the partners and families of adolescents during the pandemic (Moltrecht

Table 3
Simple logistic regression model for total sample (N = 82): Association between group to which the mother belongs, sociodemographic, obstetric, and psychosocial factors and PPD (EPDS ≥ 9)

	OR	p	95 CI
Teenage mother group			
AM-BP	Ref.	-	-
AM-DP	5.874	.001	2.263-15.348
Sociodemographic data			
Age of teenage mother	.756	.113	.535-1.068
Socioeconomic status			
A/B and C+ (High / upper-middle)	Ref.	-	-
C and C - (Middle/lower-middle)	1.071	.907	.338-3.393
D+ D and E (Low level/poverty)	2.671	.117	.782-9.122
Educational attainment			
Basic (elementary and junior high school)	1.635	.270	.683
Middle school (senior high school)	Ref.	-	-
Attended school during pregnancy			
Yes	Ref.	-	-
No	1.138	.773	.473-2.736
Attended school postpartum			
Yes	Ref.	-	-
No	1.444	.466	.537-3.884
Marital status			
Single/Separated	1.629	.313	.632-4.201
Married/Cohabiting	Ref.	-	-
In a relationship with the baby's father			
Yes	Ref.	-	-
No	1.484	.445	.539-4.087
Work (at the time of the survey)			
Yes	Ref.	-	-
No	.469	.259	.126-1.745
Obstetrics			
GW at delivery	.959	.759	.733-1.254
Birthweight (grs)	1.00	.937	.999-1.001
Psychosocial			
Maternal efficacy	.414	.001	.281-.608
Self-esteem	.753	.001	.666-.850
Dissatisfaction with social support			
Overall or total [†]	1.483	.001	1.197-1.838
Partner	1.489	.006	1.119-1.981
Family	1.785	.013	1.130-2.822
Friends	1.294	.070	.979-1.711
Depression in pregnancy			
Yes	11.393	.001	3.874-33.503
No	Ref.	-	-
Anxiety during pregnancy			
Yes	3.215	.013	1.280-8.078
No	Ref.	-	-
Planned pregnancy			
Yes	Ref.	-	-
No	1.257	.644	.477-3.314
Wanted pregnancy			
Yes	Ref.	-	-
No	1.818	.213	.710-4.659

Notes: OR = (odds ratio/odds ratio); CI = Confidence interval; AM-BP = Teenage mothers before the pandemic; AM-DP = Teenage mothers during the pandemic; GW = Gestation weeks at time of delivery.

[†]The higher the score, the greater the dissatisfaction with the social support received.

Table 4
Multiple logistic regression model for the total sample ($N = 82$): Association of group to which mother belongs and psychosocial factors with PPD ($EPDS \geq 9$)

	OR-A	p	95 CI
Group to which the teenage mother belongs			
AM-BP	Ref.	-	-
AM-DP	1.474	.613	.328 - 6.625
Psychosocial aspects			
Maternal efficacy	.663	.103	.405 - 1.087
Self-esteem	.826	.017	.737 - .971
Dissatisfaction with social support	1.430	.073	.967 - 2.13
Depression in pregnancy			
Yes	4.059	.082	.836 - 19.713
No	Ref.	-	-
Anxiety during pregnancy			
Yes	2.478	.219	.584 - 10.517
No	Ref.	-	-

Notes: OR-A = (adjusted odds ratio); CI = Confidence interval; AM-BP = Adolescent mothers before the pandemic; AM-DP = Adolescent mothers during the pandemic.
Model $X^2 = 55.995$; $p = 0.001$.

et al., 2022; Merriman et al., 2023), which impacted their perception of maternal efficacy due to their feeling that they lacked sufficient practical support and information to promote their babies' development (Moltrecht et al., 2022).

Evidence was found of higher percentages of perceived depression (82.9%) and anxiety (70%) during pregnancy among AM-DP than AM-BP. Among the former, percentages were also higher than among other groups of pregnant adolescents who experienced the pandemic: depression = 43% (Tele et al., 2022) and anxiety = 45.5% (Chamdimba et al., 2022). The perception of depression and anxiety in pregnancy of AM-DP may be overestimated since they were retrospectively evaluated with single items rather than with conventional scales as in the comparison studies (Tele et al., 2022; Chamdimba et al., 2022). It should be noted, however, that these measurements have proved effective in identifying depression and anxiety symptoms in pregnancy and postpartum (Lara et al., 2017; Patiño, 2016; Marcos-Nájera et al., 2021).

In regard to the third objective, determining whether belonging to the AM-DP group was a significant factor in experiencing PPD ($EPDS \geq 9$), simple association analyses showed that being an AM-DP, presenting with lower levels of self-esteem, greater dissatisfaction with overall social support, from their partners and families, as well as experiencing depression and anxiety during pregnancy were associated with PPD ($EPDS \geq 9$). As in previous studies, these data confirm the association between low maternal efficacy (Lara et al., 2017; Léniz-Maturana et al., 2022), low self-esteem (Birkeland et al., 2005; Ramos-Marcuse et al., 2009), dissatisfaction with social support (deCastro et al., 2011; Kim et al., 2014), having depression or anxiety during preg-

nancy (Patiño, 2016), and becoming a mother during the pandemic (Bower et al., 2023) and PPD ($EPDS \geq 9$).

A key finding was that, when controlling for the previous factors in a multiple analysis, the only significant factor for presenting with PPD ($EPDS \geq 9$) was having lower self-esteem. These data place self-esteem at the center of the relationship between psychosocial factors and PPD. There are two possible explanations for this result. First, adolescence sees a decline in self-esteem—particularly in girls—attributable to changes in body image, the emerging ability to think abstractly about oneself and the future, and to experiencing more challenging situations than in childhood (Robins & Trzesniewski, 2005). Second, being a mother can influence how adolescents perceive themselves (Mora-Guerrero et al., 2021). It can contribute to their self-esteem and identity if they regard it as an opportunity to mature and give meaning to their life, but if they regard it more as an event entailing unwanted responsibilities, it can have negative effects on their self-esteem and identity (Melgar, 2015; Patiño, 2016; Mora-Guerrero et al., 2021). The continuous discrepancy between the ideal and the real self leads to negative thoughts and feelings (Escalante, 2004), a central factor in depression (Abdel-Khalek, 2016).

This suggests that, under normal circumstances and particularly in emergencies, self-esteem should be regarded as a key element in programs to prevent PPD that can begin during pregnancy and continue in the first year postpartum. Likewise, self-esteem should be considered in interventions to treat those who have already been identified with depression in their pregnancy and postpartum. It has already been noted that social support has positive effects on the self-esteem of adolescents and their motherhood: respectful sup-

port makes it easier for them to acquire confidence in their abilities and make more appropriate decisions about their life and parenting (Mora-Guerrero et al., 2021).

Strengths: To the best of our knowledge, this is the first study in Mexico to provide information on the psychosocial and mental health factors of adolescent mothers during the pandemic and include a comparison group prior to the health crisis. In general, there is a dearth of research exploring the mental health of adolescents during pregnancy and the postpartum period compared to those of adult mothers. In this respect, this study contributes to lending visibility to the problem. On the other hand, it is relevant that the most widely recognized self-report scale (EPDS) was used to identify those with a high probability of presenting PPD. In addition, a cut-off point was selected that has shown adequate sensitivity and specificity in primiparous and postpartum adolescent mothers (Venkatesh et al., 2014). It should be noted, however, that the EPDS is not a clinical diagnosis.

Limitations: The study is cross-sectional, which only enables levels of association to be established between variables. At the same time, the data correspond to adolescent mothers living in Mexico City, who use primary care health centers (AM-BP) and Facebook (AM-DP). These circumstances limit their generalization to other groups of adolescent mothers who do not share these characteristics. The level of commitment of participants and the social desirability bias (Singh & Sagar, 2021) may differ due to the two methods of administering the survey, although it is not known how this may be reflected. Moreover, both groups primarily comprise people interested in obtaining mental health information and support for themselves, which could be particularly true of AM-DP who answered the online survey a year after the start of the health crisis. Another limitation was that since adolescent mothers were not asked about adverse events they had experienced during the pandemic, we do not know how they may have influenced PPD (EPDS \geq 9).

In conclusion, AM-DP experienced more PPD (EPDS \geq 9) than AM-BP. The crisis impacted various psychosocial factors associated with PPD (EPDS \geq 9). The role of self-esteem in PPD (EPDS \geq 9) prevails over other factors. These data show how major crises can affect this already vulnerable population and suggests elements to be considered when implementing mental health prevention and treatment measures in this population, such as boosting self-esteem.

Funding

None.

Conflict of interest

The authors declare that they have no conflicts of interest.

Acknowledgements

Thanks are due to CONAHCYT and the Master's and Doctoral Program in Medical, Dental and Health Sciences at the Universidad Nacional Autónoma de México (UNAM) for their support.

The first author received a doctoral grant with CVU number 625151. We would also like to thank María Elena Sánchez Enciso and Sarahit Guzmán Corona for collaborating with the data collection at the health centers.

REFERENCES

- Abdel-Khalek, A. M. (2016). *Introduction to the Psychology of Self-Esteem*. In F. Holloway. *Self-esteem: Perspectives, Influences, and Improvement Strategies* (pp. 1-23). New York: Nova Science Publishers.
- Alvarado-Esquivel, C., Sifuentes-Álvarez, A., & Salas-Martínez, C. (2014). The use of the Edinburgh Postpartum Depression Scale in a Population of Teenager Pregnant Women in Mexico: A Validation Study. *Clinical Practice and Epidemiology in Mental Health*, 10, 129-132. doi: 10.2174/1745017901410010129
- Alvarez-Nieto, C., Pastor-Moreno, G., Linares-Abad, M., Serrano-Martos, J., & Rodríguez-Olalla, L. (2014). Maternidad temprana: percepciones e implicaciones de las madres de las adolescentes. *Matronas Profesión*, 15(3), 88-94.
- Asociación Mexicana de Agencias de Inteligencia de Mercado y Opinión. Comité de Nivel Socioeconómico [AMAI]. (2020). *Inteligencia Aplicada a Decisiones - Julio*. Retrieved from <https://www.amai.org/NSE/index.php?queVeo=queEs> y <https://www.amai.org/NSE/index.php?queVeo=niveles> Access date April 2022
- Astle, S., Duncan, J., Toews, M., Perez-Brena, N., McAllister, P., Maddy, M., & Feinberg, M. (2021). "A Little Bit Closer": A Mixed Method Analysis of the Effect of the COVID-19 Pandemic on the Lives of Adolescent Parents. *Journal of Adolescent Research*, 1-34. doi: 10.1177/07435584211062116
- Bandura, A. (1994). Self-efficacy. In V. Ramachandran. *Encyclopedia of human behavior* (Vol. 4, pp. 71-81). New York: Academic Press.
- Barassi, F., & Grealish, A. (2022). Validity of the Edinburgh Postnatal Depression Scale for screening pregnant and postpartum adolescents: a systematic review. *Australian Journal of Advanced Nursing*, 39(2), 65-75. doi: 10.37464/2020.392.446
- Beck, C. (2001). Predictors of postpartum depression: an update. *Nursing Research*, 50(5), 275-285.
- Birkeland, R., Thompson, J., & Phares, V. (2005). Adolescent motherhood and postpartum depression. *Journal of Clinical Child and Adolescent Psychology*, 34(2), 292-300. doi: 10.1207/s15374424jccp3402_8
- Borg Xuereb, C., Borg Xuereb, R., & Jomeen, J. (2023). Adolescent Pregnancy and Early Parenting. In R. Borg Xuereb, & J. Jomeen. *Perspectives on Midwifery and Parenthood*. Cham, Switzerland: Springer International Publishing.
- Bower, M., Smout, S., Donohoe-Bales, A., O'Dean, S., Teesson, L., Boyle, J., Lim, D., Nguyen, A., Callear, A., Batterham, P., Gournay, K., & Teesson, M. (2023). A hidden pandemic? An umbrella review of global evidence on mental health in the time of COVID-19. *Frontiers in Psychiatry*, 14, 1107560. doi: 10.3389/fpsyt.2023.1107560
- Burki, T. (2020). The indirect impact of COVID-19 on women. *The Lancet*, 20(8), 904-905. doi: 10.1016/S1473-3099(20)30568-5
- Chamdimba, E., Munthali, A., Kabiru, C., Thakwalakwa, C., Ushie, B., & Ajayi, A. (2022). Self-report of mental health distress among pregnant and parenting adolescents during the COVID-19 pandemic in Malawi. PREPRINT available at *Research Square*. doi: 10.21203/rs.3.rs-1809318/v1
- Cox, J., Holden, J., & Sagovsky, R. (1987). Detection of Postnatal Depression: Development of the 10-item Edinburgh Postnatal Depression Scale. *British Journal of Psychiatry*, 150(6), 782-786. doi: 10.1192/bjp.150.6.782
- deCastro, F., Hinojosa-Ayala, N., & Hernandez-Prado, B. (2011). Risk and protective factors associated with postnatal depression in Mexican adolescents. *Journal of Psychosomatic Obstetrics & Gynecology*, 32(4), 210-217. doi: 10.3109/0167482X.2011.626543
- Edward, M., & Mobarak, A. (2022). The economics of the COVID-19 pandemic in poor countries. *Annual Review of Economics*, 14(1), 253-285. doi: 10.3386/w29339
- Escalante, G. (2004). *Autoestima y diferenciación personal*. Universidad de los Andes, Centro de Investigaciones Psicológicas, Mérida, Venezuela.
- Farkas-Klein, C. (2008). Escala de evaluación parental (EEP): desarrollo, propiedades psicométricas y aplicaciones. *Universitas Psychologica*, 7(2), 457-467.
- Ganchimeg, T., Ota, E., Morisaki, N., Laopaiboon, M., Lumbiganon, P., Zhang, J., Yamdamsuren, B., Temmerman, M., Say, L., Tunçalp, Ö., Vogel, J. P., Souza, J.

- P., Mori, R., & WHO Multicountry Survey on Maternal Newborn Heal. (2014). Pregnancy and childbirth outcomes among adolescent mothers: a World Health Organization multicountry study. *BJOG: An International Journal of Obstetrics & Gynaecology*, 121(S1), 40-48.
- Gniewosz, G. (2022). A mother's perspective: perceived stress and parental self-efficacy during the COVID-19 pandemic. *European Journal of Developmental Psychology*, 20(4), 666-693. doi: 10.1080/17405629.2022.2120464
- Goto, R., Piedvache, A., Hangai, M., Yamaoka, Y., Sampei, M., Sawada, N., Okubo, Y., Tanaka, K., Morisaki, N., & Hosozawa, M. (2022). Time trends in emotional well-being and self-esteem in children and adolescents during the COVID-19 pandemic. *Child and Adolescent Psychiatry and Mental Health*, 16(89), 2-10. doi: 10.1186/s13034-022-00525-3
- Hodgkinson, S., Beers, L., Southammakosane, C., & Lewin, A. (2014). Addressing the Mental Health Needs of Pregnant and Parenting Adolescents. *Pediatrics*, 133(1), 114-122. doi: 10.1542/peds.2013-0927
- Ibarra-Yruegas, B., Lara, M. A., Navarrete, L., Nieto, L., & Kawas Valle, O. (2016). Psychometric properties of the Postpartum Depression Predictors Inventory-Revised for pregnant women in Mexico. *Journal of Health Psychology*, 23(11), 1415-1423. doi: 10.1177/1359105316658969
- International Business Machines. (2017). *IBM SPSS Statistics for Macintosh, Version 25.0*. Armonk, NY: IBM Corp.
- Kim, T., Connolly, J., & Tamim, H. (2014). The effect of social support around pregnancy on postpartum depression among Canadian teen mothers and adult mothers in the maternity experiences survey. *BMC Pregnancy and Childbirth*, 14(1), 162.
- Lara, M. A., Patiño, P., Navarrete, L., Hernández, Z., & Nieto, L. (2017). Association between depressive symptoms and psychosocial factors and perception of maternal self-efficacy in teenage mothers. *Salud Mental*, 40(5), 201-208. doi: 10.17711/SM.0185-3325.2017.026
- Lara, M., Navarrete, L., Medina, E., Patiño, P., & Tiburcio, M. (2023). Impact of Facebook on Social Support and Emotional Wellbeing in Perinatal Women during Three Waves of the COVID-19 Pandemic in Mexico: A Descriptive Qualitative Study. *International Journal of Environmental Research and Public Health*, 20(3), 2472. doi: 10.3390/ijerph20032472
- Lazarus, R., & Folkman, S. (1987). Transactional theory and research on emotions and coping. *European Journal of Personality*, 1(3), 141-169. doi: 10.1002/per.2410010304
- Léniz-Maturana, L., Vilaseca, R., & Leiva, D. (2022). Maternal self-efficacy and emotional well-being in Chilean adolescent mothers: the relationship with their children's social-emotional development. *PeerJ*, 10, e13162. doi: 10.7717/peerj.13162
- López-Romo, H. (2020). *Cátedra Prima Webinar: Los mexicanos a través de una mirada a los Niveles Socioeconómicos* – Agosto 28, 2020. Retrieved from <https://youtu.be/Dr5wb81Vfdc> Access date Abril 2022
- Marcos-Nájera, R., Rodríguez-Muñoz, M. F., Lara, M. A., Navarrete, L., & Le, H. N. (2021). A cross-cultural analysis of the prevalence and risk factors for prenatal depression in Spain and Mexico. *Culture, Medicine, and Psychiatry*, 45(4), 599-612.
- Matei, A., Dimitriu, M., Cirstoveanu, C., Socea, B., & Ionescu, C. (2021). Assessment of Postpartum Depression in Adolescents Who Delivered during COVID-19 Social Restrictions: The Experience of a Tertiary Hospital from Bucharest, Romania. *Healthcare*, 9(7), 807. doi: 10.3390/healthcare9070807
- Medina-Jimenez, V., Bermudez-Rojas, M., Murillo-Vargas, H., Rivera-Camarillo, A., Muñoz-Acosta, J., Ramirez-Abarca, T., Esparza-Valencia, D., Angeles-Torres, A., Lara-Avila, L., Hernandez-Muñoz, V., Madrigal-Tejeda, F., Estudillo-Jimenez, G., Jacobo-Enciso, L., Torres-Torres, J., Espino-y-Sosa, S., Baltazar-Martinez, M., Villanueva-Calleja, J., Nava-Sanchez, A., Mendoza-Carrera, C., ... Martinez-Portilla, R. (2022). The impact of the COVID-19 pandemic on depression and stress levels in pregnant women: A national survey during the COVID-19 pandemic in Mexico. *The Journal of Maternal-Fetal & Neonatal Medicine*, 35(23), 4438-4441. doi: 10.1080/14767058.2020.1851675
- Melgar, L. (2015). Adolescentes traicionadas: Derecho a la salud, el bienestar de las jóvenes y las omisiones del Estado. En M. Tapia, I. Saucedo, & L. Ramos. *Violencia de género, juventud y escuelas en México* (pp. 137-144). México: Centro de estudios para el Adelanto de las Mujeres y la Equidad de Género.
- Merriman, B., Jarmoc, G., van der Rijn, M., & Pierre-Joseph, N. (2023). Impact of COVID-19 on Mental Health and Resiliency of Pregnant and Parenting Adolescents and Young Adults: A Qualitative Study. *Journal of Pediatric Health Care*, 37(5), 484-491. doi: 10.1016/j.pedhc.2023.03.002
- Moltrecht, B., Dalton, L., Hanna, J., Law, C., & Rapa, E. (2022). Young parents' experiences of pregnancy and parenting during the COVID-19 pandemic: a qualitative study in the United Kingdom. *BMC Public Health*, 22(1), 523. doi: 10.1186/s12889-022-12892-9
- Mora-Guerrero, G., Escárate-Colín, L., Espinoza-Lerdón, C., & Peña-Paredes, A. (2021). Apoyo social percibido, autoestima y maternidad adolescente: entre el respeto y la intrusión. Estudio en Traiguén, Chile. *Prospectiva - Revista de Trabajo Social e Intervención Social*, (32), 151-171.
- NU - CEPAL. (2020). *Risks of the COVID-19 pandemic for the exercise of women's sexual and reproductive rights*. Retrieved from <https://repositorio.cepal.org/handle/11362/46508> Access date April 2023
- Patiño, P. (2016). *Factores de riesgo, síntomas de depresión y ansiedad en el posparto en adolescentes* [Tesis de Maestría en Ciencias de la Salud]. México: Universidad Nacional Autónoma de México. Retrieved from <http://132.248.9.195/ptd2016/noviembre/0753731/0753731.pdf>
- Ramos-Marcuse, F., Oberlander, F., Papas, F., McNary, S., Hurley, K., & Black, M. (2009). Stability of maternal depressive symptoms among urban, low-income, African American adolescent mothers. *Journal of Affective Disorders*, 122(1-2), 68-75. doi: 10.1016/j.jad.2009.06.018
- Records, K., Rice, M., & Beck, C. (2007). Psychometric assessment of the Postpartum Depression Predictors Inventory-Revised. *Journal of Nursing Measurement*, 15(3), 189-202. doi: 10.1891/106137407783095775
- Rivera-Rivera, N. Y., McGuinn, L., Osorio-Valencia, E., Martínez-Medina, S., Schnaas, L., Wright, R. J., Téllez-Rojo, M. M., Wright, R. O., Tamayo-Ortiz, M., & Rosa, M. J. (2021). Changes in Depressive Symptoms, Stress and Social Support in Mexican Women during the COVID-19 Pandemic. *International Journal of Environmental Research and Public Health*, 18(16), 8775. doi: 10.3390/ijerph18168775
- Robins, R., & Trzesniewski, K. (2005). Self-Esteem Development Across the Life Span. *Current Directions in Psychological Science*, 14(3), 158-162. doi: 10.1111/j.0963-7214.2005.00353.x
- Rosenberg, M. (1965). *Society and the adolescent self-image*. Princeton, New Jersey: Princeton University Press.
- Sánchez-Tanquer, M., González-Pier, E., Sepúlveda, J., Abascal-Miguel, L., & Fieldhouse, J. (2021). *La respuesta de México al Covid-19: Estudio de caso*. Retrieved from <https://globalhealthsciences.ucsf.edu/news/mexicos-response-covid-19-case-study> Access date March 2023
- Sangsawang, N., & Sangsawang, B. (2023). Postpartum depression, social support and maternal self-efficacy between adolescent and adult mothers during the COVID-19 pandemic: A comparative cross-sectional study. *Journal of Advanced Nursing*, 79(1), 113-124. doi: 10.1111/jan.15445
- Shigemura, J., Ursano, R., Morganstein, J., Kurosawa, M., & Benedek, D. (2020). Public responses to the novel 2019 coronavirus (2019-nCoV) in Japan: Mental health consequences and target populations. *Psychiatry and Clinical Neurosciences*, 74(4), 281-282. doi: 10.1111/pcn.12988
- Singh, S., & Sagar, R. (2021). A critical look at online survey or questionnaire-based research studies during COVID-19. *Asian Journal of Psychiatry*, 65, 102850. doi: 10.1016/j.ajp.2021.102850
- Smiley, Y., Sadeghi, N., Jolda, C., & Chokshi, B. (2021). Parenting in a Pandemic: Needs of Teen Parents During COVID-19. *Clinical Pediatrics*, 60(14), 559-563. doi: 10.1177/00099228211054296
- Tele, A., Kathono, J., Mwaniga, S., Nyongesa, V., Yator, O., Gachuno, O., Wamalwa, D., Amugune, B., Cuijpers, P., Saxena, S., McKay, M., Carvajal, L., Lai, J., Huang, K. Y., Merali, Z., & Kumar, M. (2022). Prevalence and risk factors associated with depression in pregnant adolescents in Nairobi, Kenya. *Journal of Affective Disorders Reports*, 10, 100424. doi: 10.1016/j.jadr.2022.100424
- UNFPA. (2020). *Consecuencias socioeconómicas del embarazo en la adolescencia en seis países de América Latina y el Caribe. Implementación de la Metodología Milena en Argentina, Colombia, Ecuador, Guatemala, México y Paraguay*. Fondo de Población de las Naciones Unidas - Oficina Regional.
- UNFPA. (2023). *El Fondo de Población de las Naciones Unidas (México) Abril 7, 2023*. Retrieved from <https://mexico.unfpa.org/es/news/conapo-imjuve>

el-fondo-de-poblaci%C3%B3n-de-las-naciones-unidas-unfpa-y-organon-impulsan-un-proyecto#:~:text=En%20su%20intervenci%C3%B3n%2C%20la%20secretaria,m%C3%A1s%20de%20mil%20por%20d%C3%ADa Access date April 2023

Venkatesh, K., Zlotnick, C., Triche, E., Ware, C., & Phipps, M. (2014) Accuracy of brief screening tools for identifying postpartum depression among adolescent mothers. *Pediatrics*, 133(1), e45-e53. doi: 10.1542/peds.2013-1628

World Health Organization [WHO]. (2022). *The impact of COVID-19 on mental health cannot be made light of* – June 16, 2022. Retrieved from World Health

Organization: <https://www.who.int/news-room/feature-stories/detail/the-impact-of-covid-19-on-mental-health-cannot-be-made-light-of#:~:text=A%20great%20number%20of%20people,affected%20much%20more%20than%20others> Access date April 2023

Xue, A., Oros, V., La Marca-Ghaemmaghami, P., Scholkmann, F., Righini-Grunder, F., Natalucci, G., Karen, T., Bassler, D., & Restin, T. (2021). New Parents Experienced Lower Parenting Self-Efficacy during the COVID-19 Pandemic Lockdown. *Children*, 8(2), 79. doi: 10.3390/children8020079